



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/44, 31/24 C07D 213/40, C07C 69/616	A1	(11) International Publication Number: WO 92/17176 (43) International Publication Date: 15 October 1992 (15.10.92)
(21) International Application Number: PCT/US92/02120 (22) International Filing Date: 13 March 1992 (13.03.92) (30) Priority data: 675,780 27 March 1991 (27.03.91) US (71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, One Abbott Park Road, Abbott Park, IL 60064-3500 (US). (72) Inventors: SHAM, Hing, L. ; 5109 Pembroke Court, Gurnee, IL 60031 (US). KEMPF, Dale, J. ; 2565 Tyler Court, Libertyville, IL 60048 (US). NORBECK, Daniel, W. ; 307 Hazelwood Drive, Lindenhurst, IL 60047 (US). ZHAO, Chen ; 5225 Frederick Court, Gurnee, IL 60031 (US).		(74) Agents: GORMAN, Edward, Hoover, Jr. et al.; Abbott Laboratories, CHAD 0377/AP6D-2, One Abbott Park Road, Abbott Park, IL 60064-3500 (US). (81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), MC (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: RETROVIRAL PROTEASE INHIBITING COMPOUNDS (57) Abstract Compounds are disclosed which are retroviral protease inhibitors. Also disclosed are methods of using the compounds and compositions for inhibiting a retroviral protease and for treating an HIV infection.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
DE	Germany	MC	Monaco	TG	Togo
DK	Denmark			US	United States of America

RETROVIRAL PROTEASE INHIBITING COMPOUNDS

This is a continuation-in-part of U.S. patent application Serial No. 518,730, filed May 9, 1990, which is a continuation-in-part of U.S. patent application Serial No. 456,124, filed December 22, 1989, which is a continuation-in-part of U.S. patent application Serial No. 405,604, filed September 8, 1989, which is a continuation-in-part of U.S. patent application Serial No. 355,945, filed May 23, 1989.

Technical Field

This invention was made with Government support under contract number AI27220-01 awarded by the National Institute of Allergy and Infectious Diseases. The Government has certain rights in this invention.

The present invention relates to novel compounds and a composition and method for inhibiting retroviral proteases and in particular for inhibiting human immunodeficiency virus (HIV) protease, a composition and method for treating

-2-

a retroviral infection and in particular an HIV infection, processes for making such compounds and synthetic intermediates employed in these processes.

Background Art

Retroviruses are those viruses which utilize a ribonucleic acid (RNA) intermediate and a RNA-dependent deoxyribonucleic acid (DNA) polymerase, reverse transcriptase, during their life cycle. Retroviruses include, but are not limited to, the RNA viruses of the Retroviridae family, and also the DNA viruses of the Hepadnavirus and Caulimovirus families. Retroviruses cause a variety of disease states in man, animals and plants. Some of the more important retroviruses from a pathological standpoint include human immunodeficiency viruses (HIV-1 and HIV-2), which cause acquired immune deficiency syndrome (AIDS) in man, hepatitis B virus, which causes hepatitis and hepatic carcinomas in man, human T-cell lymphotropic viruses I, II, IV and V, which cause human acute cell leukemia, and bovine and feline leukemia viruses which cause leukemia in domestic animals.

Proteases are enzymes which cleave proteins at specific peptide bonds. Many biological functions are controlled or mediated by proteases and their complementary protease inhibitors. For example, the protease renin cleaves the peptide angiotensinogen to produce the peptide angiotensin I. Angiotensin I is further cleaved by the protease angiotensin converting enzyme (ACE) to form the hypotensive peptide angiotensin II. Inhibitors of renin and ACE are known to reduce high blood pressure in vivo. An inhibitor of a retroviral

-3-

protease should provide a therapeutic agent for diseases caused by the retrovirus.

The genomes of retroviruses encode a protease that is responsible for the proteolytic processing of one or more polyprotein precursors such as the pol and gag gene products. See Wellink, Arch. Virol. 98 1 (1988). Retroviral proteases most commonly process the gag precursor into core proteins, and also process the pol precursor into reverse transcriptase and retroviral protease. In addition, retroviral proteases are sequence specific. See Pearl, Nature 328 482 (1987).

The correct processing of the precursor polyproteins by the retroviral protease is necessary for the assembly of infectious virions. It has been shown that in vitro mutagenesis that produces protease-defective virus leads to the production of immature core forms which lack infectivity. See Crawford, J. Virol. 53 899 (1985); Kato, et al., Virology 145 280 (1985). Therefore, retroviral protease inhibition provides an attractive target for antiviral therapy. See Mitsuya, Nature 325 775 (1987).


Current treatments for viral diseases usually involve administration of compounds that inhibit viral DNA synthesis. Current treatments for AIDS (Dagani, Chem. Eng. News, November 23, 1987 pp. 41-49) involve administration of compounds such as 2',3'-dideoxycytidine, trisodium phosphonoformate, ammonium 21-tungsto-9-antimoniate, 1-beta-D-ribofuranosyl-1,2,4-triazole-3-carboxamide, 3'-azido-3'-deoxythymidine, and adriamycin that inhibit viral DNA synthesis; compounds such as AL-721 and polymannoacetate which may prevent HIV from

-4-

penetrating the host cell; and compounds which treat the opportunistic infections caused by the immunosuppression resulting from HIV infection. None of the current AIDS treatments have proven to be totally effective in treating and/or reversing the disease. In addition, many of the compounds currently used to treat AIDS cause adverse side effects including low platelet count, renal toxicity and bone marrow cytopenia.


Disclosure of the Invention

In accordance with the present invention, there are retroviral protease inhibiting compounds of the formula:



wherein R₁ is loweralkyl, cycloalkyl, phenyl or substituted derivatives thereof wherein the loweralkyl, cycloalkyl or phenyl group is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo;
R₂ is phenyl, aromatic heterocyclic or substituted phenyl wherein the phenyl ring is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo;

-5-

R₃ is

wherein m is 0 to 3 and R₄ is independently selected at each occurrence from hydrogen, loweralkyl, cycloalkylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, thioalkoxyalkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl and aminocarbonylalkyl and



wherein R₆ is aryl, heterocyclic or substituted aryl wherein the aryl ring is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo, V is O or S, W is CH₂, -CH₂CH₂-, -CH=CH- or -D-CH=CH- wherein D is O, S or NH and Y is absent, O, S or N(R₇) wherein R₇ is hydrogen or loweralkyl; or T is R₁₅C(O)-, R₁₅C(S)- or R₁₅S(O)₂- wherein R₁₅ is loweralkyl, aryl or heterocyclic; and Z is -C(O)- or -CH(OH)-; or a pharmaceutically acceptable salt thereof; with the proviso that the compound is other than 5-(N-benzyloxycarbonyl-isoleucyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

The compounds of this invention having one asymmetric carbon atom can exist as the pure enantiomers or as the racemic mixture. The compounds of this invention having two or more asymmetric carbon atoms can exist as pure

-6-

diastereomers, mixtures of diastereomers, diastereomeric racemates or mixtures of diastereomeric racemates. The present invention includes within its scope all of the isomeric forms. The terms "S" and "R" configuration as used herein are as defined by the IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, Pure Appl. Chem. (1976) 45, 13-30.

The terms "Gly" and "Val" as used herein refer to glycine and valine, respectively. In general, the amino acid abbreviations used herein follow the IUPAC-IUB Joint Commission on Biochemical Nomenclature for amino acids and peptides (Eur. J. Biochem. 1984, 158, 9-31).

The term "loweralkyl" as used herein refers to straight or branched chain alkyl radicals containing from 1 to 6 carbon atoms including, but not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, n-pentyl, 1-methylbutyl, 2,2-dimethylbutyl, 2-methylpentyl, 2,2-dimethylpropyl, n-hexyl and the like.

The term "cycloalkyl" as used herein refers to an aliphatic ring having 3 to 7 carbon atoms including, but not limited to, cyclopropyl, cyclopentyl, cyclohexyl and the like.

The term "aryl" as used herein refers to a monocyclic or bicyclic carbocyclic ring system having one or more aromatic rings including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl and the like.

The term "arylalkyl" as used herein refers to an aryl group appended to a loweralkyl radical including, but not limited to, benzyl, 1-naphthylmethyl and the like.

The term "alkylamino" as used herein refers to a loweralkyl radical appended to an NH radical.

-7-

The term "dialkylamino" as used herein refers to $-NR_{30}R_{31}$ wherein R_{30} and R_{31} are independently selected from loweralkyl groups.

The term "heterocyclic ring" or "heterocyclic" as used herein refers to any 3- or 4-membered ring containing a heteroatom selected from oxygen, nitrogen and sulfur; or a 5- or 6-membered ring containing one, two or three nitrogen atoms; one nitrogen and one sulfur atom; or one nitrogen and one oxygen atom. The 5-membered ring has 0-2 double bonds and the 6-membered ring has 0-3 double bonds. The nitrogen heteroatoms can be optionally quaternized. The term "heterocyclic" also includes bicyclic groups in which any of the above heterocyclic rings is fused to a benzene ring or a cyclohexane ring or another heterocyclic ring. Heterocyclics include: pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, imidazolyl, imidalinyl, imidazolidinyl, pyridyl, piperidinyl, pyrazinyl, piperazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzothiazolyl, benzoxazolyl, furyl, thienyl and benzothienyl.

Heterocyclics can be unsubstituted or monosubstituted or disubstituted with substituents independently selected from hydroxy, halo, oxo ($=O$), alkylimino ($R^*N=$ wherein R^* is a loweralkyl group), amino, alkylamino, dialkylamino, alkoxy, polyalkoxy, haloalkyl, cycloalkyl, aryl, arylalkyl, $-COOH$, $-SO_3H$ and loweralkyl. In addition, nitrogen containing heterocycles can be N-protected.

-8-

The term "halo" or "halogen" as used herein refers to -Cl, -Br, -I or -F.

The terms "alkoxy" and "thioalkoxy" as used herein refer to $R_{32}O-$ and $R_{32}S-$, respectively, wherein R_{32} is a loweralkyl group or benzyl.

The term "polyalkoxy" as used herein refers to $-OR_{33}$ wherein R_{33} is a straight or branched chain containing 1-5, C_n-O-C_n linkages wherein n' and n'' are independently selected from 1 to 3, including but not limited to methoxyethoxymethoxy, methoxymethoxy and the like.

The term "haloalkyl" as used herein refers to a loweralkyl radical in which one or more of the hydrogen atoms are replaced by halogen including, but not limited to, chloromethyl, trifluoromethyl, 1-chloro-2-fluoroethyl and the like.

The term "cycloalkylalkyl" as used herein refers to a loweralkyl radical to which is appended a cycloalkyl group including, but not limited to, cyclohexylmethyl and the like.

The term "hydroxyalkyl" as used herein refers to a loweralkyl radical to which is appended a hydroxy group (-OH).

The term "alkoxyalkyl" as used herein refers to a loweralkyl radical to which is appended an alkoxy group.

The term "thioalkoxyalkyl" as used herein refers to a loweralkyl radical to which is appended an alkoxy group.

The term "aminoalkyl" as used herein refers to a loweralkyl radical to which is appended an amino group ($-NH_2$).

-9-

The term "alkylaminoalkyl" as used herein refers to a loweralkyl radical to which is appended an alkylamino group (i.e., $R_{40}NH-$ wherein R_{40} is loweralkyl).

The term "dialkylaminoalkyl" as used herein refers to a loweralkyl radical to which is appended a dialkylamino group (i.e., $R_{41}R_{42}N-$ wherein R_{41} and R_{42} are independently selected from loweralkyl).

The term "carboxyalkyl" as used herein refers to a loweralkyl radical to which is appended a carboxyl group ($-COOH$).

The term "alkoxycarbonylalkyl" as used herein refers to a loweralkyl radical to which is appended an alkoxycarbonyl group (i.e., $R_{43}OC(O)-$ wherein R_{43} is loweralkyl).

The term "aminocarbonylalkyl" as used herein refers to a loweralkyl radical to which is appended an aminocarbonyl group (i.e., $-C(O)NH_2$).

The term "N-protected" or "N-protecting group" as used herein refers to those groups intended to protect the N-terminus of an amino acid or peptide or to protect an amino group against undesirable reactions during synthetic procedures. N-protecting groups comprise carbamates, amides, N-alkyl derivatives, amino acetal derivatives, N-benzyl derivatives, imine derivatives, enamine derivatives and N-heteroatom derivatives, all of which are disclosed in Greene, "Protective Groups In Organic Synthesis," John Wiley & Sons, New York (1981), which is incorporated by reference herein. In particular, N-protecting groups include formyl, benzoyl, pivaloyl, benzyl, phenylsulfonyl, t-butyloxycarbonyl (Boc), benzyloxycarbonyl (Cbz) and the like.

-10-

Representative compounds of the invention include;

5(S) - (N-benzyloxycarbonyl-(L)-isoleucyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-benzyloxycarbonyl-(L)-tert-leucyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-benzyloxycarbonyl-(L)-asparaginy1)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-Benzyloxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(2-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(3-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(4-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(2-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(3-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(4-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(2-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(3-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(4-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S) - (N-(2-(2-Pyridyl)ethanesulfonyl)-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;

-11-

5(S)-(N-Acetylglycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane; and
5(S)-(N-(Methanesulfonyl)-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.


The compounds of the invention can be prepared as outlined in the processes shown in Schemes I and II. According to Scheme I, oxazolidinone **1** (wherein R_1 is defined as above and R_{10} is an activating group, such as $-N(OCH_3)(CH_3)$ and the like) is reacted with an organometallic reagent R_2CH_2M (**2**) (wherein R_2 is defined as above and M is a metal ion, for example, R_2CH_2Li , R_2CH_2MgCl and the like). Oxazolidinone **1** can be prepared according to the procedure disclosed in European Patent Application No. EP402646, published December 19, 1990 (U.S. patent application Serial No. 518,730, filed May 9, 1990, which is incorporated by reference herein). Difluoroketone **3** is then converted to **4** (for example, by reduction to the alcohol, formation of the mesylate, elimination to the olefin and hydrogenation). Deprotection of **4** (for example, by treatment with barium hydroxide and the like), followed by coupling with carboxylic acid **5**, or an activated derivative thereof, provides difluoroalcohol **6a**. Oxidation of **6a** (for example, with $Na_2Cr_2O_7$ /acetic acid and the like) provides **6b**.

Scheme II illustrates an alternative process for the preparation of the deprotected form of **4** (i.e., **12**). N-protected amino alcohol **7** (R_1 is as defined above and R_{20} is an N-protecting group, for example,


-12-

t-butyloxycarbonyl) is oxidized (for example, by Swern oxidation) to provide aldehyde 8. Reaction of aldehyde 8 with substituted acetylene 9 (R_2 is defined as above) and zinc in an inert solvent provides difluoro alcohol 10. Hydrogenation of 10 gives 11, which can be deprotected to provide 12.

-13-


SCHEME I

-14-

SCHEME II

-15-

Intermediates useful for the preparation of compounds I include compounds of the formula:



wherein R_1 is loweralkyl, cycloalkyl, phenyl or substituted derivatives thereof wherein the loweralkyl, cycloalkyl or phenyl group is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo;

R_2 is phenyl, aromatic heterocyclic or substituted phenyl wherein the phenyl ring is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo;

R_{21} is hydrogen or an N-protecting group; and

R_{22} is hydrogen, or R_{21} and R_{22} taken together are $-C(O)-$.

Other useful intermediates for the preparation of compounds I include compounds of the formula:



-16-

wherein m is 0 to 3 and R₄ is independently selected at each occurrence from hydrogen, loweralkyl, cycloalkylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, thioalkoxyalkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl and aminocarbonylalkyl and



wherein R₆ is aryl, heterocyclic or substituted aryl wherein the aryl ring is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo, V is O or S, W is CH₂, -CH₂CH₂-, -CH=CH- or -D-CH=CH- wherein D is O, S or NH and Y is absent, O, S or N(R₇) wherein R₇ is hydrogen or loweralkyl; or T is R₁₅C(O)-, R₁₅C(S)- or R₁₅S(O)₂- wherein R₁₅ is loweralkyl, aryl or heterocyclic; or an acid halide or activated ester derivative thereof .

Acid halide derivatives of the above intermediates include the acid chloride. Activated ester derivatives of the above intermediates include activated esters commonly used by those skilled in the art for activating carboxylic acid groups for coupling with an amine to form a peptide bond including, but not limited to, formic and acetic acid derived anhydrides, anhydrides derived from alkoxycarbonyl halides such as isobutyloxycarbonylchloride and the like, N-hydroxysuccinimide derived esters, N-hydroxyphthalimide derived esters, N-hydroxybenzotriazole derived esters, N-hydroxy-5-norbornene-2,3-dicarboxamide derived esters,

-17-

4-nitrophenol derived esters, 2,4,5-trichlorophenol derived esters and the like.

The following examples will serve to further illustrate the preparation of the novel compounds of the invention.

Example 1

5(S)-(N-Benzylloxycarbonyl-(L)-valinyl)-amino-1-(2-pyridyl)-6-phenyl-3,3-difluoro-4-oxo-hexane

1A. 4(S)-Benzyl-5(R)-(2'-(2',2'-difluoro-N'-methoxy-N'-methylacetamido)-2-oxazolidinone

To a solution of 1.36 g of the 2-oxazolidinone derivative of 4(S)-amino-5-phenyl-2,2-difluoro-3(R)-hydroxypentanoic acid (European Patent Application No. EP402646, published December 19, 1990) in 20 ml of dimethylformamide (DMF) was added successively, 615 mg of N,O-dimethyl hydroxylamine hydrochloride, 2.11 ml of triethylamine and 1.47 g of 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride. The reaction mixture was stirred at room temperature overnight. The solvent was removed in vacuo and the residue was extracted with ethyl acetate and washed successively with saturated aqueous sodium carbonate and saturated aqueous sodium chloride, then dried and concentrated. Silica gel chromatography (20% ethyl acetate/methylene chloride) provided 1.3 g of the desired product. ^1H NMR (CDCl_3): δ 2.85 (m, 2H), 3.10 (m, 2H), 3.20 (s, 3H), 3.73 (s, 3H), 4.10 (m, 1H), 4.90 (m, 1H), 5.10 (br s, 1H), 7.25 (m, 5H).

-18-

1B. 4(S)-Benzyl-5(R)-(3'-(3',3'-difluoro-2'-oxo-1'-(2-pyridyl)propyl))-2-oxazolidinone

To 500 mg of the product from Example 1A in 10 ml of tetrahydrofuran (THF) at -78°C was added a solution of the lithium derivative of 2-picoline (generated by adding 2.5 ml of a 1.6 M solution of n-BuLi in hexane to 0.39 ml of 2-picoline in 5 ml of THF at -78°C). After stirring for 30 minutes at -78°C, and then for 30 minutes at -10°C, the reaction mixture was quenched by adding saturated aqueous ammonium chloride solution. This mixture was then extracted with ethyl acetate. The combined organic layer was dried and concentrated. Silica gel chromatography (2-5% methanol/methylene chloride) provided 405 mg of the desired product. ¹H NMR (CDCl₃): δ 2.80 (m, 1H), 3.05 (m, 1H), 4.30 (m, 1H), 4.85 (m, 1H), 5.0 (br s, 1H), 5.90 (s, 1H), 7.05-7.30 (m, 7H), 7.70 (m, 1H), 8.05 (d, 1H).

1C. 5(S)-amino-1-(2-pyridyl)-6-phenyl-3,3-difluoro-4(R)-hydroxyhexane

To 150 mg of the product from Example 1B in methanol at 0°C was added a solution of 30 mg of sodium borohydride in 10 ml of methanol. After stirring for 30 minutes, the product was isolated. The crude product was treated with 1.1 equivalents of triethylamine and 1.0 equivalents of methanesulfonyl chloride in methylene chloride to provide the mesylate derivative. The crude mesylate was treated with 5 equivalents of DBU in 5 ml of toluene at 80°C to provide the olefin. The olefin was hydrogenated using 10% Pd/C as catalyst and methanol as solvent. The resulting product was hydrolyzed using barium hydroxide in

-19-

1:1 dioxane/water to provide the desired product. Mass spectrum: (M+1)=307.

1D. 5(S)-(N-Benzylloxycarbonyl-(L)-valinyl)-amino-1-(2-pyridyl)-6-phenyl-3,3-difluoro-4-oxo-hexane

To 100 mg of the resultant product of Example 1C in 5 ml of DMF was added successively, 100 mg of N-benzylloxycarbonyl-(L)-valine, 160 mg of HOBt, 0.117 ml of triethylamine and 160 mg of 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride. The reaction mixture was stirred at room temperature overnight. The product was isolated and purified by silica gel chromatography (2% methanol/methylene chloride) to provide 145 mg of the crude alcohol, which was oxidized by dissolving in 5 ml of acetic acid and adding 1.1 equivalents of sodium dichromate (Synthesis, 466(1989)), resulting in the desired product. ¹H NMR (CDCl₃): δ 0.81 (d, 3H), 0.90 (d, 3H), 2.05 (m, 1H), 2.50-3.30 (m, 6H), 3.95 (m, 1H), 5.05 (s, 2H), 5.20 (br d, 1H), 5.30 (m, 1H), 6.30 (br d, 1H), 7.10-7.35 (m, 7H), 7.60 (m, 1H), 8.50 (m, 1H). Mass spectrum: (M+1)=538.

Example 2.

5(S)-(N-(2-Pyridyl)-methoxycarbonyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

4(S)-Benzyl-5(R)-(3'-(3',3'-difluoro-1'-phenylpropyl)-2-oxazolidinone (European Patent Application No. EP402646, published December 19, 1990) was deprotected by treatment with barium hydroxide. The resultant difluoroamino alcohol was coupled to 5-(N-(2-

-20-

pyridyl)methoxycarbonyl)valine using N-ethyl-N'-(dimethylaminopropyl)carbodiimide, 1-hydroxybenzotriazole and trimethylamine in dimethylformamide (DMF). Oxidation of the resultant compound according to Example 1D using sodium dichromate in acetic acid provided the desired compound (65%). ^1H NMR (CDCl_3): δ 0.82 (d, 3H), 0.90 (d, 3H), 2.05 (m, 1H), 2.35 (m, 2H), 2.80-2.95 (m, 2H), 3.30 (m, 1H), 3.95 (q, 1H), 5.20 (s, 2H), 5.30 (m, 1H), 6.20 (brd, 1H), 7.10-7.30 (m, 12H), 7.70 (t, 1H), 8.60 (d, 1H).

Example 3

5(S)-(N-(2-Pyridyl)-methylamino-carbonyl-(L)-Valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

Using the procedure of Example 2, but replacing 5-(N-(2-pyridyl)-methoxycarbonyl)-valine with 5-(N-(2-pyridyl)-methylamino-carbonyl)-Valine provided the desired compound. ^1H NMR (CDCl_3): δ 0.82 (d, 3H), 0.90 (d, 3H), 2.07 (m, 1H), 2.30 (m, 1H), 2.80-3.30 (m, 4H), 4.10 (q, 1H), 4.45 (d, 1H), 5.30 (m, 1H), 5.55 (t, 1H), 6.50 (d, 1H), 7.12-7.30 (m, 12H), 7.65 (m, 1H), 8.50 (m, 1H).

Example 4

5(S)-(N-Benzylloxycarbonyl-glycyl-(L)-Valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

Using the procedure in Example 2, but replacing 5-(N-(2-pyridyl)methoxycarbonyl)-valine with 5-N-Cbz-glycyl-valine provided the desired product. ^1H NMR (CDCl_3): δ 0.80 (d, 3H), 0.88 (d, 3H), 2.05 (m, 1H), 2.40 (m, 2H),

-21-

2.70-3.30 (m, 4H), 3.70 (m, 2H), 4.20 (q, 1H), 5.10 (s, 2H), 5.30 (m, 1H), 6.30 (m, 2H), 7.12-7.38 (m, 15H).

Example 5

N-t-Butyloxycarbonyl-(1,6-diphenyl-3,3-difluoro-4(R,S)-hydroxy-5(S)-amino-hexyne

To a solution of 1 gm of Boc-phenylalaninal in 12 ml of THF was added 1.05 gm of zinc dust and 0.16 gm of mercuric chloride. To this sonicated suspension was added a solution of 0.25 gm of bromodifluoromethylphenylacetylene in 4 ml of THF over a one hour period. Excess zinc was filtered and the filtrate concentrated and washed with 10% aq. KHSO₄ solution. Extraction with EtOAc (3 x 100 ml), drying with anhydrous Na₂SO₄ and evaporation of the solvent in vacuo provided a yellow oil which was purified by silica gel column chromatography to give 1.09 gm of the desired product (68%). Mass spectrum: (M + 1) = 402. ¹H NMR (CDCl₃): δ 1.34 (s, 9H), 2.95 - 3.20 (m, 2H), 3.90 - 4.10 (m, 2H), 5.05 (br d, 1H), 7.25 - 7.50 (m, 10H).

Example 6

N-t-butyloxycarbonyl-(1,6-diphenyl-3,3-difluoro-4(R,S)-hydroxy-5(S)-amino-hexane).

To a suspension of 21 mg of 10% palladium on carbon was added a methanol solution of the resultant product from Example 5. The suspension was stirred vigorously under a hydrogen atmosphere. After one hour, the catalyst was filtered off and

-22-

the filtrate concentrated to provide 81 mg (95%) of desired product. Mass spectrum: $(M + 1) = 406$. ^1H NMR (CDCl_3): δ 1.40 (s, 9H), 2.25 (m, 2H), 2.70 - 3.10 (m, 4H), 3.50 (m, 1H), 3.70 (m, 1H), 4.0 (m, 1H), 4.90 (br d, 1H), 7.25 (m, 10H).

Example 7

1,6-Diphenyl-3,3-difluoro-4(R,S)-hydroxy-5(S)-amino-hexane.

To 1 gm of the resultant product from Example 6 was added 5 ml of trifluoroacetic acid and 5 ml of dichloromethane. After one hour at room temperature, the solution was concentrated to an oil. The oil was dissolved in 100 ml EtOAc and washed with saturated NaHCO_3 solution. The organic layer was dried with anhydrous Na_2SO_4 and evaporation of the solvent in vacuo provided a white solid, 740 mg (98%). Mass spectrum: $(M + 1) = 306$.

Example 8

5(S)-(N-(2-(2-Pyridyl)ethanesulfonyl)-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

Coupling of N-(2-(2-pyridyl)ethanesulfonyl)-glycyl-valine to the resultant product from Example 7 using the procedure of Example 2, followed by oxidation with sodium dichromate in acetic acid provided the desired product. Mass spectrum: $(M + 1) = 629$. ^1H NMR (CDCl_3): δ 0.75 (d, 3H), 0.85 (d, 3H), 2.10 (m, 1H),

-23-

2.40 (m, 2H), 2.80 (m, 3H), 3.30 (m, 3H), 3.60 (m, 2H), 3.68 (br d, 2H), 4.20 (m, 1H), 5.30 (m, 1H), 5.95 (m, 1H), 6.60 (d, 1H), 6.80 (d, 1H), 7.20 (m, 12H), 7.60 (m, 1H), 8.50 (d, 1H).

Example 9

5(S)-(N-Acetylglycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

Using the procedure described in Example 8, but replacing N-(2-(2-pyridylethane)sulfonyl)-glycyl-valine with N-acetyl-glycyl-valine provided the desired product. Mass spectrum: (M + 1) = 502. ¹H NMR (CDCl₃): δ 0.80 (d, 3H), 0.88 (d, 3H), 2.03 (s, 3H), 2.02 (m, 1H), 2.45 (m, 2H), 2.80 (t, 2H), 2.92 (m, 1H), 3.30 (m, 1H), 3.89 (m, 2H), 4.20 (m, 1H), 5.30 (m, 1H), 6.20 (m, 1H), 6.40 (t, 2H), 7.25 (m, 10H).

Example 10

5(S)-(N-(Methanesulfonyl)-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

Using the procedure described in Example 8, but replacing N-(2-(2-pyridyl)ethanesulfonyl)-glycyl-valine with N-(methanesulfonyl)-glycyl-valine provided the desired product. Mass spectrum: (M + 1) = 538. ¹H NMR (CDCl₃): 0.81 (d, 3H), 0.88 (d, 3H), 2.05 (m, 1H), 2.35 (m, 2H), 2.85 (m, 3H), 2.95 (s, 3H), 3.30 (dd, 1H), 3.80 (d, 2H), 4.20 (m, 1H), 5.20 (m, 1H), 5.30 (m, 1H), 6.30 (d, 1H), 6.55 (d, 1H), 7.25 (m, 10H).

-24-

Example 11

Using the methods outlined above, the following compounds can be prepared:

- (a) 5-(N-benzyloxycarbonyl-(L)-isoleucyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (b) 5-(N-benzyloxycarbonyl-(L)-tert-leucyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (c) 5-(N-benzyloxycarbonyl-(L)-asparaginyll)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (d) 5(S)-(N-(2-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (e) 5(S)-(N-(3-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (f) 5(S)-(N-(4-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (g) 5(S)-(N-(2-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (h) 5(S)-(N-(3-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (i) 5(S)-(N-(4-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (j) 5(S)-(N-(2-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
- (k) 5(S)-(N-(3-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane; and
- (l) 5(S)-(N-(4-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

-25-

Fluorogenic Assay for Screening Inhibitors of HIV
Protease

The inhibitory potency of the compounds of the invention can be determined by the following method.

A compound of the invention is dissolved in DMSO and a small aliquot further diluted with DMSO to 100 times the final concentration desired for testing. The reaction is carried out in a 6 X 50 mm tube in a total volume of 300 microliters. The final concentrations of the components in the reaction buffer are: 125 mM sodium acetate, 1 M sodium chloride, 5 mM dithiothreitol, 0.5 mg/ml bovine serum albumin, 1.3 μ M fluorogenic substrate, 2% (v/v) dimethylsulfoxide, pH 4.5. After addition of inhibitor, the reaction mixture is placed in the fluorometer cell holder and incubated at 30°C for several minutes. The reaction is initiated by the addition of a small aliquot of cold HIV protease. The fluorescence intensity (excitation 340 nM, emission 490 nM) is recorded as a function of time. The reaction rate is determined for the first six to eight minutes. The observed rate is directly proportional to the moles of substrate cleaved per unit time. The IC₅₀ is the concentration of test compound needed to inhibit the protease by 50%.

Fluorogenic substrate: Dabcyl-Ser-Gln-Asn-Tyr-Pro-Ile-Val-Gln-EDANS wherein DABCYL = 4-(4-dimethylamino-phenyl)azobenzoic acid and EDANS = 5-((2-aminoethyl)amino)-naphthalene-1-sulfonic acid.

Table 1 shows the inhibitory potencies of compounds of the invention against HIV-1 protease.

-26-

TABLE 1

<u>Compound of Example</u>	<u>IC₅₀ (nM)</u>
1	14
2	2.2
3	2.9
4	1.0
8	2.5
9	3.2
10	2.2
11c	21

The results indicate that the compounds are effective as inhibitors of HIV-1 protease.

Antiviral Activity

The anti-HIV activity of the compounds of the invention can be determined in MT4 cells according to the procedure of Pauwels et. al. (*J. Virol. Methods* 1988, 20, 309). The IC₅₀ is the concentration of compound that gives 50% inhibition of the cytopathic effect of HIV. The LC₅₀ is the concentration of compound at which 50% of the cells remain viable.

Table 2 shows the inhibitory potencies of compounds of the invention against HIV-1_{3B} in MT4 cells.

-27-

Table 2

<u>Compound of Example</u>	<u>IC₅₀ (μM)</u>	<u>LC₅₀ (μM)</u>
1	5.7	>100
2	6.3	21
3	7.6	>100
4	1.9	>100
8	4.9	>100
9	8.6	>100
10	4.3	>100
11c	10.4	>100

The results indicate that the compounds are effective as inhibitors of the cytopathic effect of HIV in cell culture.

The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as loweralkyl halides,

-28-

such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid. Other salts include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or with organic bases.

The compounds of the invention are useful for inhibiting a retroviral protease, in particular HIV protease, in vitro or in vivo. The compounds of the present invention are also useful for the treatment or prophylaxis of diseases caused by retroviruses, especially acquired immune deficiency syndrome or an HIV infection in a human or other mammal.

Total daily dose administered to a host in single or divided doses may be in amounts, for example, from 0.001 to 100 mg/kg body weight daily and more usually 0.1 to 10 mg. Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

-29-

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination, and the severity of the particular disease undergoing therapy.

The compounds of the present invention may be administered orally, parenterally, sublingually, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Injectable preparations, for example, sterile injectable aqueous or oleagenous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-propanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including

-30-

synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

The compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or

-31-

multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and phosphatidyl cholines (lecithins), both natural and synthetic.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

While a compound of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or more immunomodulators, antiviral agents, other anti-infective agents or vaccines. Other antiviral agents to be administered in combination with a compound of the present invention include AL-721, beta interferon, polymannoacetate, ganciclovir, dideoxycytidine, trisodium phosphonoformate, HPA-23, eflonithine, Peptide T, Reticulose (nucleophosphoprotein), zidovudine (AZT), ansamycin LM 427, trimetrexate, UA001, ribavirin, alpha interferon, oxetanocin, DDC, DDI, cylobut G and acyclovir. Immunomodulators that can be administered in combination with a compound of the present invention include bropirimine, Ampligen, anti-human alpha interferon antibody, colony stimulating factor, CL246,738, Imreg-1, Imreg-2, diethyldithiocarbamate, interleukin-2, alpha-interferon, inosine pranobex, methionine enkephalin, muramyl-tripeptide, TP-5, erythropoietin, naltrexone and

-32-

tumor necrosis facator. Other antiinfective agents that can be administered in combination with a compound of the present invention include pentamidine isethionate. Any of a variety of HIV or AIDS vaccines can be used in combination with a compound of the present invention.

It will be understood that agents which can be combined with the compounds of the present invention for the treatment or prophylaxis of AIDS or an HIV infection are not limited to those listed above, but include in principle any agents useful for the treatment or prophylaxis of AIDS or an HIV infection.

When administered as a combination, the therapeutic agents can be formulated as separate compositions which are given at the same time or different times, or the therapeutic agents can be given as a single composition.


The foregoing is merely illustrative of the invention and is not intended to limit the invention to the disclosed compounds. Variations and changes which are obvious to one skilled in the art are intended to be within the scope and nature of the invention which are defined in the appended claims.

-33-

CLAIMS

What is claimed is:


1. A compound of the formula:



wherein R_1 is loweralkyl, cycloalkyl, phenyl or substituted derivatives thereof wherein the loweralkyl, cycloalkyl or phenyl group is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo;

R_2 is phenyl, aromatic heterocyclic or substituted phenyl wherein the phenyl ring is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo;

R_3 is



wherein m is 0 to 3 and R_4 is independently selected at each occurrence from hydrogen, loweralkyl, cycloalkylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, thioalkoxyalkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl and aminocarbonylalkyl and

-34-




wherein R_6 is aryl, heterocyclic or substituted aryl wherein the aryl ring is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo, V is O or S, W is CH_2 , $-\text{CH}_2\text{CH}_2-$, $-\text{CH}=\text{CH}-$ or $-\text{D}-\text{CH}=\text{CH}-$ wherein D is O, S or NH and Y is absent, O, S or N(R_7) wherein R_7 is hydrogen or loweralkyl; or T is $\text{R}_{15}\text{C}(\text{O})-$, $\text{R}_{15}\text{C}(\text{S})-$ or $\text{R}_{15}\text{S}(\text{O})_2-$ wherein R_{15} is loweralkyl, aryl or heterocyclic; and Z is $-\text{C}(\text{O})-$ or $-\text{CH}(\text{OH})-$; or a pharmaceutically acceptable salt thereof; with the proviso that the compound is other than 5-(N-benzyloxycarbonyl-isoleucyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane.

2. The compound of Claim 1 wherein R_1 is phenyl; R_2 is phenyl, 2-pyridyl, 3-pyridyl or 4-pyridyl; m is 2; R_4 is independently selected at each occurrence from hydrogen and isopropyl; Z is $-\text{C}(\text{O})-$ and T is benzyloxycarbonyl, N-(2-pyridylmethyl)aminocarbonyl, N-(3-pyridylmethyl)aminocarbonyl, N-(4-pyridylmethyl)aminocarbonyl, 2-pyridylmethoxycarbonyl, 3-pyridylmethoxycarbonyl, 4-pyridylmethoxycarbonyl, N-(2-pyridylmethyl)-N-(methyl)aminocarbonyl, N-(3-pyridylmethyl)-N-(methyl)aminocarbonyl or N-(4-pyridylmethyl)-N-(methyl)aminocarbonyl.

-35-


3. A compound of the formula:



wherein R₁ is phenyl;

R₂ is phenyl;

R₃ is



wherein m is 2, R₄ is independently selected at each occurrence from hydrogen and loweralkyl and




wherein R₆ is aryl, heterocyclic or substituted derivatives thereof wherein the aryl or heterocyclic ring is substituted with one or two substituents independently selected from hydroxy, alkoxy, thioalkoxy and halo, V is O or S, W is CH₂, -CH₂CH₂-, -CH=CH- or -D-CH=CH- wherein D is O, S or NH and Y is absent, O, S or N(R₇) wherein R₇ is hydrogen or loweralkyl; or T is R₁₅C(O)-, R₁₅C(S)- or R₁₅S(O)₂- wherein R₁₅ is loweralkyl, aryl or heterocyclic; or a pharmaceutically acceptable salt thereof.

-36-

4. The compound of Claim 3 wherein R_4 is independently selected at each occurrence from hydrogen and isopropyl and T is benzyloxycarbonyl, N-(2-pyridylmethyl)aminocarbonyl, N-(3-pyridylmethyl)aminocarbonyl, N-(4-pyridylmethyl)aminocarbonyl, 2-pyridylmethoxycarbonyl, 3-pyridylmethoxycarbonyl, 4-pyridylmethoxycarbonyl, N-(2-pyridylmethyl)-N-(methyl)-aminocarbonyl, N-(3-pyridylmethyl)-N-(methyl)-aminocarbonyl or N-(4-pyridylmethyl)-N-(methyl)-aminocarbonyl.


5. A compound of the formula:



wherein R_1 is phenyl;

R_2 is phenyl;

R_3 is



wherein R_4 is isopropyl and



T is

wherein R_6 is aryl, heterocyclic or substituted derivatives thereof wherein the aryl or heterocyclic ring is substituted with one or two substituents independently selected from

-37-

hydroxy, alkoxy, thioalkoxy and halo, V is O or S, W is CH₂, -CH₂CH₂-, -CH=CH- or -D-CH=CH- wherein D is O, S or NH and Y is absent, O, S or N(R₇) wherein R₇ is hydrogen or loweralkyl; or T is R₁₅C(O)-, R₁₅C(S)- or R₁₅S(O)₂- wherein R₁₅ is loweralkyl, aryl or heterocyclic; or a pharmaceutically acceptable salt thereof.

6. The compound of Claim 7 wherein T is benzyloxycarbonyl, N-(2-pyridylmethyl)aminocarbonyl, N-(3-pyridylmethyl)aminocarbonyl, N-(4-pyridylmethyl)aminocarbonyl, 2-pyridylmethoxycarbonyl, 3-pyridylmethoxycarbonyl, 4-pyridylmethoxycarbonyl, N-(2-pyridylmethyl)-N-(methyl)-aminocarbonyl, N-(3-pyridylmethyl)-N-(methyl)-aminocarbonyl or N-(4-pyridylmethyl)-N-(methyl)-aminocarbonyl.

7. A compound selected from the group consisting of:
5(S)-(N-Benzyloxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S)-(N-(2-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S)-(N-(3-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S)-(N-(4-pyridyl)methoxycarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S)-(N-(2-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S)-(N-(3-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
5(S)-(N-(4-pyridylmethyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;


-38-

5(S) - (N-(2-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
 5(S) - (N-(3-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
 5(S) - (N-(4-pyridylmethyl)-N-(methyl)aminocarbonyl-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
 5(S) - (N-(2-(2-Pyridyl)ethanesulfonyl)-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane;
 5(S) - (N-Acetylglycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane; and
 5(S) - (N-(Methanesulfonyl)-glycyl-(L)-valinyl)-amino-3,3-difluoro-1,6-diphenyl-4-oxo-hexane; or a pharmaceutically acceptable salt thereof.

8. A method for inhibiting a retroviral protease comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.


9. A method for treating an HIV infection comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.

10. A process for the preparation of a compound of Claim 1 comprising reacting a compound of the formula:




-39-

with a compound of the formula:



INTERNATIONAL SEARCH REPORT

International Application No. PCT/US92/02120

I. CLASSIFICATION F SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC ⁵ A61K 31/44, 31/24; C07D 213/40, C07C 69/616 U.S. Cl. 546/332, 335; 560/27; 514/357, 539		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
US	546/332, 335; 514/357, 539; 560/27	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	Chemical abstract, vol. 110, No. 19, issued May 8, 1989 (Columbus, Ohio USA) (Schirlin et al.) Preparation of activated electrophilic ketone tetroamide analogs of peptidase substrates as peptidase inhibitors, abstract no. 173757V.	6,7
A	Chemical abstract, vol. 106, No. 11, issued March, 1987 (Columbus, Ohio USA) amino acid and peptide derivatives as peptidase inhibitors, abstract no. 85059F.	6,7
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁴ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
16 June 1992		3 0 JUN 1992
International Searching Authority		Signature of Authorized Officer
ISA/US		 Jan T. Fan

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter ¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹³, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).


VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This International Searching Authority found multiple inventions in this international application as follows:

See Attached Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

Claims 1-10 wherein T =  - W - Y - C^V

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
☐ No protest accompanied the payment of additional search fees.